



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MECHANICS.

NOTE.—Prize Problem No. 86, is open for solution as only one solution has been received, and that one is faulty.

93. Proposed by WALTER H. DRANE, Graduate Student, Harvard University, Cambridge, Mass.

A small rope, which is passed over a smooth pulley, has attached at one end a weight of twenty pounds, and at the other end hangs a monkey, also weighing twenty pounds. Is it possible for the monkey to climb to the pulley, and if so, what will happen to the weight?

Solution by the PROPOSER.

When a body is in motion, its weight is altered, becoming greater when the body moves upward, and less when the body moves downward, and this increase or decrease in weight is proportional to the velocity. By this principle, when the monkey starts upward with a velocity, $ds/dt = \frac{1}{2} Ft$, where F is the force with which he raises himself, his weight is increased by an amount proportional to $\frac{1}{2} Ft$. This force reacts undiminished upon the weight (as the pulley is smooth) and as the mass of the weight is also 20 pounds it must also begin to ascend with the same velocity as the monkey. This in turn increases the weight of the weight until it equals that of the monkey, which counteracts the effect of the monkey's increase in weight, and hence we have the result that monkey and weight will ascend with equal velocities.

94. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

In a parallelogram $ABCD$, $\angle D = \beta$, $AB = a$, $BC = b$, the principal moments of inertia at the centroid are $(\frac{1}{24}m)[a^2 + b^2 \pm \sqrt{(a^4 - b^4 + 2a^2b^2\cos 2\beta)}]$ and the principal axes at the same point make with the side CD an angle θ given by

$$\tan 2\theta = \frac{b^2 \sin 2\beta}{a^2 + b^2 \cos 2\beta}.$$

Solution by the PROPOSER.

Let $ABCD$ be the parallelogram, O the intersection of the diagonals AC , BD , and OX , OY be the axes. For integration we will transform to OX , OY .

$$\tan 2\theta = \frac{2 \sum mxy}{\sum mx^2 - \sum my^2}.$$

$\sum mx^2$ = moment of inertia about OX

$$= \rho \sin \beta \int_{-\frac{1}{2}a}^{\frac{1}{2}a} \int_{-\frac{1}{2}b}^{\frac{1}{2}b} (x + y \cos \beta)^2 dx dy = \frac{1}{2} \rho a b \sin \beta (a^2 + b^2 \cos^2 \beta) = \frac{1}{2} m (a^2 + b^2 \cos^2 \beta).$$

$$\sum my^2 = \rho \sin^3 \beta \int_{-\frac{1}{2}a}^{\frac{1}{2}a} \int_{-\frac{1}{2}b}^{\frac{1}{2}b} y^2 dx dy = \frac{1}{2} m b^2 \sin^2 \beta.$$